

- reservoir: A comparison with the external catheter. *JPEN* 13:478-483, 1989.
22. Brothers TE, Von Moll LK, Niederhuber JE, et al.: Experience with subcutaneous infusion ports in three hundred patients. *Surg Gynecol Obstet* 166:295-301, 1988.
 23. Alastrué A, Rull M, Escudero LE, et al.: Experiencia con 150 reservorios venosos subcutáneos como acceso venoso e infusión de tratamiento en pacientes adultos oncológicos, hematológicos y afectados de síndrome de inmunodeficiencia adquirida. *Med Clin (Barc)* 99:444-449, 1992.
 24. Wisborg T, Flaatten H, Koller M-E: Percutaneous placement of permanent central venous catheters: experience with 200 catheters. *Acta Anaesthesiol Scand* 35:49-51, 1991.
 25. Carles J, Rosell R, Ribelles N, et al.: Sistema de acceso venoso con reservorio subcutáneo en la población oncológica. Experiencia en 60 pacientes. *Oncología* 14:196-199, 1991.
 26. Abraham JL, Mullen JL: A prospective study of prolonged central venous access in leukemia. *JAMA* 248:2868-2873, 1982.
 27. Pagelow C, Narvaez M, Toledano S, et al.: Experience with a totally implantable venous device in children. *Am J Dis Child* 140:69-71, 1986.
 28. Ryan JA, Abel RM, Abbott WM, et al.: Catheter complications in total parenteral nutrition. *N Engl J Med* 290:757-761, 1974.
 29. Murphy IM, Lipman TO: Central venous catheter care in parenteral nutrition: A review. *JPEN* 11:190-201, 1987.
 30. Reed WP, Newman KA, Applefeld MM, et al.: Drug extravasation as a complication of venous access ports. *Ann Intern Med* 102:788-789, 1985.
 31. Sariago J, Bootorabi B, Matsumoto T, et al.: Major long-term complications in 1422 permanent venous access devices. *Am J Surg* 165:249-251, 1993.

COMMENTARY

This interesting paper demonstrates the utility of the external jugular vein for the passage of a long-term intra-

venous catheter connected to a totally implanted device for long-term chemotherapy.

In the United States, the cephalic vein was used initially for cannulation through a cut-down in the deltopectoral groove. However, in a minority of patients this vein was absent or of insufficient size to be cannulated. Occasionally, although the cephalic vein was capacious enough to allow its cannulation, the catheter could not be advanced past the cephalic-subclavian vein junction. Similar experiences were noted in a few instances when the external jugular vein was used by this reviewer for cannulation. In some patients, the angle of the junction of external jugular or cephalic vein with the subclavian is such that the catheter as it goes through tends to hit the opposite wall of the vein and not to advance centrally.

The most popular route in the United States is currently the subclavian vein through direct cannulation, which is easy and safe with the available kits for insertion, including the peel-away catheters. This paper, however, indicates that with experience the external jugular vein can be an effective alternative for insertion of these catheters.

Constantine P. Karakousis, MD, PhD

Millard Fillmore Hospital and
State University of New York at Buffalo
Buffalo, New York 14209